

Percent of births to mothers who smoked during pregnancy in each racial/ethnic group, 2012-2016

A healthy start in life begins during pregnancy and maternal smoking diminishes that good start. Non-Hispanic White Kentuckians are more likely to smoke during pregnancy than other racial/ethnic groups. Characteristics affecting the likelihood of smoking while pregnant include: marital status, socioeconomic status, access to prenatal care, and living in a racially segregated area.

	Black (non-Hispanic)	Hispanic	White (non-Hispanic)
Kentucky	14.3%	4.3%	22.8%
Boone	7.0%	2.6%	16.5%
Campbell	20.6%	5.6%	21.0%
Christian	12.7%	5.3%	18.0%
Daviess	17.9%	S	17.6%
Fayette	16.4%	2.1%	13.2%
Hardin	10.2%	11.6%	19.6%
Jefferson	11.5%	3.3%	14.1%
Kenton	26.8%	4.9%	23.3%
McCracken	13.2%	10.9%	18.0%
Warren	13.1%	2.0%	15.0%

S = Data is suppressed for fewer than 6 births.

Source: Kentucky Cabinet for Health and Family Services, Vital Statistics Branch, processed by the Kentucky State Data Center.

Data note: Data in highlighted cells should be used with caution as rates were calculated with relatively small numbers.

Percent of births to mothers who smoked during pregnancy in each racial/ethnic group, 2011-2015

A healthy start in life begins during pregnancy and maternal smoking diminishes that good start. Non-Hispanic White Kentuckians are more likely to smoke during pregnancy than other racial/ethnic groups. Characteristics affecting the likelihood of smoking while pregnant include: marital status, socioeconomic status, access to prenatal care, and living in a racially segregated area.

	Black (non-Hispanic)	Hispanic	White (non-Hispanic)
Kentucky	14.9%	4.3%	23.6%
Boone	6.9%	2.5%	17.3%
Campbell	21.3%	6.3%	21.4%
Christian	12.8%	4.6%	18.9%
Daviess	20.8%	S	18.9%
Fayette	17.1%	2.2%	13.8%
Hardin	10.3%	10.4%	20.1%
Jefferson	12.2%	3.3%	14.6%
Kenton	28.8%	5.7%	23.7%
McCracken	14.0%	9.6%	20.2%
Warren	14.6%	2.2%	16.4%

S = Data is suppressed for fewer than 6 births.

Source: Kentucky Cabinet for Health and Family Services, Vital Statistics Branch, processed by the Kentucky State Data Center.

Data note: Data in highlighted cells should be used with caution as rates were calculated with relatively small numbers.

Percent of births to mothers who smoked during pregnancy in each racial/ethnic group, 2010-2014

A healthy start in life begins during pregnancy and maternal smoking diminishes that good start. Non-Hispanic White Kentuckians are more likely to smoke during pregnancy than other racial/ethnic groups. Characteristics affecting the likelihood of smoking while pregnant include: marital status, socioeconomic status, access to prenatal care, and living in a racially segregated area.

	Black (non-Hispanic)	Hispanic	White (non-Hispanic)
Kentucky	15.6%	4.3%	24.2%
Boone	7.1%	2.1%	17.6%
Campbell	22.8%	4.8%	22.1%
Christian	14.1%	5.1%	19.9%
Daviess	22.1%	S	21.0%
Fayette	17.7%	2.2%	14.1%
Hardin	11.5%	9.7%	20.2%
Jefferson	13.0%	3.3%	15.4%
Kenton	27.9%	5.7%	24.3%
McCracken	14.7%	9.7%	20.6%
Warren	14.8%	2.2%	18.0%

S = Data is suppressed for fewer than 6 births.

Source: Kentucky Cabinet for Health and Family Services, Vital Statistics Branch, processed by the Kentucky State Data Center.

Data note: Data in highlighted cells should be used with caution as rates were calculated with relatively small numbers.

Percent of births to mothers who smoked during pregnancy in each racial/ethnic group, 2009-2013

A healthy start in life begins during pregnancy and maternal smoking diminishes that good start. Non-Hispanic White Kentuckians are more likely to smoke during pregnancy than other racial/ethnic groups. Characteristics affecting the likelihood of smoking while pregnant include: marital status, socioeconomic status, access to prenatal care, and living in a racially segregated area.

	Black (non-Hispanic)	Hispanic	White (non-Hispanic)
Kentucky	16.2%	4.3%	24.9%
Boone	8.0%	2.9%	18.7%
Campbell	21.6%	4.6%	24.1%
Christian	14.7%	5.6%	21.7%
Daviess	22.0%	S	21.7%
Fayette	18.4%	2.0%	14.9%
Hardin	12.9%	9.5%	20.6%
Jefferson	13.4%	3.2%	16.3%
Kenton	30.1%	5.7%	25.4%
McCracken	16.7%	10.9%	22.9%
Warren	15.7%	1.8%	18.8%

S = Data is suppressed for fewer than 6 births.

Source: Kentucky Cabinet for Health and Family Services, Vital Statistics Branch, processed by the Kentucky State Data Center.

Data note: Data in highlighted cells should be used with caution as rates were calculated with relatively small numbers.